

February 1, 2008

LBNL-50258
UCB-PTH-02/21
hep-ph/0205111

Neutrino mixing and large CP violation in B physics *

Darwin Chang^{1,2}, Antonio Masiero³ and Hitoshi Murayama^{1,4}

¹*Theoretical Physics Group*

*Ernest Orlando Lawrence Berkeley National Laboratory
University of California, Berkeley, California 94720, USA*

²*Physics Department and NCTS,
National Tsing-Hua University, Hsinchu, 30043 Taiwan, ROC*

³*Dip. di Fisica "G. Galilei", Univ. di Padova and
INFN, Sez. di Padova, Via Marzolo, 8, I-35131 Padua, Italy*

⁴*Department of Physics, University of California
Berkeley, California 94720, USA*

Abstract

We show that in see-saw models of neutrino mass à la SUSY $SO(10)$, the observed large mixing in atmospheric neutrinos naturally leads to large $b - s$ transitions. If the associated new CP phase turns out to be large, this SUSY contributions can drastically affect the CP violation in some of the B decay channels yielding the β and γ angles of the unitarity triangle. They can even produce sizeable CP asymmetries in some decay modes which are not CP violating in the standard model context. Hence the observed large neutrino mixing makes observations of low energy SUSY effect in some CP violating decay channels potentially promising in spite of the agreement between the Standard Model and data in K and B physics so far.

*This work was supported in part by the U.S. Department of Energy under Contracts DE-AC03-76SF00098, in part by the National Science Foundation under grant PHY-95-14797. HM was also supported by Alfred P. Sloan Foundation.

1 Introduction

In the last couple of years we have obtained three major pieces of information on flavor physics from experiments: large neutrino mixing in atmospheric neutrinos, existence of “direct CP violation” in the neutral kaon system ϵ'/ϵ and observation of CP violation in $B \rightarrow \psi K_s$. Only the first of these three results clearly calls for new physics beyond the standard model (SM) for its explanation. As for ϵ'/ϵ , the theoretical uncertainties surrounding the SM predictions prevent any firm conclusion. Nevertheless, it is reassuring to notice that in the most familiar and promising extension of the SM, the minimal supersymmetric SM (MSSM), there is indeed room for a large contribution to ϵ'/ϵ even in the presence of a tiny deviation from flavor universality in the terms which break SUSY softly [1]. As for the observed CP violation in B physics, the results are in agreement with the SM expectation, but they leave open the possibility of large CP violating contributions from new physics (in particular the MSSM) in other decay channels which should be observable in a near future at B factories or hadron colliders.

The new physics involved to explain the atmospheric neutrino observations must provide a mass to neutrinos and guarantee at least one near-maximal mixing in the leptonic sector. One of the best candidates to accomplish these tasks is the see-saw mechanism [2, 3]. It has been known for a long time that, at variance with what occurs in the non-SUSY case, the SUSY version of the see-saw mechanism can potentially lead to large lepton flavor violating (LFV) effects [4][†]. Obviously, then, if one combines the SUSY see-saw with the idea of some hadron-lepton unification, one may suspect that the large mixing between second and third generation in the neutrino case entails not only some large LFV, but also some large mixing among quarks of the second and third generation which sit together with the leptons in a GUT multiplet. This is indeed what may happen for the right-handed quark supermultiplets in a SUSY $SU(5)$ construction with a see-saw mechanism [6]

Motivated by the three above experimental observations and the above-mentioned theoretical considerations in the SUSY context, this Letter addresses the following question: in a SUSY see-saw context where neutrino and up-quark couplings are GUT-unified, how large can the SUSY contributions to CP violation in B physics be? So far it has been observed that, considering a model-independent parameterization of the CP violating SUSY

[†]For recent works on LFV in SUSY see-saw models, see Ref. [5]

contributions in B physics, it is possible, compatibly with all the existing phenomenological constraints, to obtain sizeable effects. However, given that we have now a precise indication of large LFV effects in the neutrino sector, we find it timely and important to link this experimental fact to predictions for B physics in a motivated SUSY GUT context which accounts for the atmospheric neutrino results.

We find that : i) differently from $SU(5)$ -like schemes where one has to assume the largeness of some neutrino coupling to infer large quark flavor violation (FV) from the observed large neutrino mixing, in a $SO(10)$ -like context the link between this latter phenomenon and large $b - s$ transitions is automatically ensured; ii) the mixing $B_s - \bar{B}_s$ can receive large SUSY contributions comparable, if not larger, than the SM contribution; iii) some of the CP violating B decays which yield the β and γ angles of the unitarity triangle are strongly affected by the presence of the SUSY CP violating contributions, whilst other decays which in the SM are predicted to yield the same angles are essentially untouched by SUSY; iv) there exist some decay channels which do not present any sizeable CP asymmetry in the pure SM which develop significant (and hopefully observable) CP signals thanks to the large SUSY contributions. The Letter shows that, differently from some pessimistic common lore after the CP and FCNC results in K and B physics so far, the important experimental finding in atmospheric neutrinos yields expectations of sizeable deviations from SM in some CP violating B decays in SUSY GUT schemes where neutrino masses arise from a see-saw mechanism.

2 Main Point

The main point of this Letter is very simple. Consider an $SO(10)$ grand-unified theory, which breaks to $SU(5)$ at, *e.g.*, 10^{17} GeV. The Yukawa coupling of “third-generation” neutrino is unified with the large top Yukawa coupling thanks to the $SO(10)$ unification. However, the large mixing angle in atmospheric neutrino oscillation suggests that this “third-generation” neutrino is actually a near-maximal mixture of ν_μ and ν_τ . On the other hand, the “third-generation” charged leptons and down-type quarks have a relatively large unified bottom and tau Yukawa coupling, which is diagonal in the $SU(5)$ multiplet that contains ν_τ by definition. Therefore, the $SU(5)$ multiplet with the large top Yukawa coupling contains approximately

$$5_3^* = 5_\tau^* \cos \theta + 5_\mu^* \sin \theta, \quad (1)$$

where $\theta \simeq 45^\circ$ is the atmospheric neutrino mixing angle, and

$$5_\tau^* = (b^c, b^c, b^c, \nu_\tau, \tau) \quad (2)$$

$$5_\mu^* = (s^c, s^c, s^c, \nu_\mu, \mu). \quad (3)$$

It is interesting that even a large mixing in right-handed down quarks does not appear in CKM matrix simply because there is no charged-current weak interaction on right-handed quarks. Therefore the mixing among right-handed quarks decouples from low-energy physics.

However, the mixing among squarks yield observable effects. The top Yukawa coupling then generates an $O(1)$ radiative correction to the mass of $\tilde{s} \sin \theta + \tilde{b} \cos \theta$, which leads to a large mixing between $\tilde{s}$ and $\tilde{b}$ at low energies. This large mixing in turn generates interesting effects in B -physics. Examples include: large B_s mixing and CP violation, different “ $\sin 2\beta$ ” in $B_d \rightarrow \phi K_s$ from that in $B_d \rightarrow J/\psi K_s$ due to CP-violating penguin contribution, and different “ γ ” values from different processes.[‡]

The rest of the paper is devoted to more details of this simple point. In particular, we demonstrate in the next section that one can write down semi-concrete models of $SO(10)$ unification which lead to large $\tilde{b}$ - $\tilde{s}$ mixing. It is important that such models do not necessarily cause too-large $\mu \rightarrow e\gamma$ or other dangerous effects. Then we will discuss detailed consequences of large $\tilde{b}$ - $\tilde{s}$ in B physics.

3 $SO(10)$

3.1 Framework

We are motivated by $SO(10)$ unification [9], and we describe our assumptions on the unified framework in this section. When $SO(10)$ is broken to $SU(5)$, we have small mixings among 10 ’s responsible for CKM mixing, while we need large mixings among $\bar{5}$ ’s to explain the MNS mixing matrix [10] in the neutrino sector. Then back in the $SO(10)$ multiplets, the top quark comes together with the near-maximal linear combination of second- and third-generation $\bar{5}$ ’s. To the extent that we ignore all small Yukawa couplings except the top Yukawa coupling, the only effect of the Yukawa coupling

[‡]In models of low-energy gauge mediation [7] or anomaly mediation [8] of supersymmetry breaking, effects of our interest are simply not present.

appears in the multiplet

$$16_{i=3} \ni (t_L, V_{tn} d_{Ln}) + (t_R)^C + U_{n3}^* (d_{Rn})^C + (\nu_{L3}, U_{n3}^* l_{Ln}) + V_{tn} (l_{Rn})^C, \quad (4)$$

where we have use indices $i, j, \cdot$ to represent the indices in Y_u^D basis while $n, m, \cdot$ are reserved for the basis in which Y_d is diagonal. Here, V is the CKM matrix, and U the MNS matrix (some additional CP violating phases will be taken care of later), given the large U_{23} as evidenced in the atmospheric neutrino data, a near-maximal linear combination of s_R and b_R experiences the large top Yukawa coupling in the $SO(10)$ theory. This simple point produces a large radiative correction to the soft mass of this linear combination, which is flavor-off-diagonal in the mass basis of down quarks. Therefore we can expect a potentially large effect in B_s mixing, and other related effects in B -physics.

Our framework is closely related to that in [11] with the superpotential

$$W = \frac{1}{2}(Y_u)_{ij} 16_i 16_j 10_u + \frac{1}{2}(Y_d)_{ij} 16_i 16_j 10_d. \quad (5)$$

Here, Y_u, Y_d are up- and down-type Yukawa matrices, $i, j = 1, 2, 3$ are generation indices, and $10_u, 10_d$ are Higgs multiplets that contain H_u and H_d in the MSSM.

In addition to the above two terms, W must include some further (renormalizable or non renormalizable) Yukawa coupling responsible for the right-handed neutrino masses. As usual, this can be achieved either through $\langle 126 \rangle$ or $\langle \bar{16} \rangle^2 / M_{Pl}$.

We need at least two Yukawa matrices Y_u and Y_d to generate intergenerational mixings and hence two Higgs multiplets. In this sense, this is the “minimal” framework of $SO(10)$ unification. However, this makes both Y_u and Y_d matrices symmetric. A symmetric Y_u is acceptable phenomenologically, while a symmetric requirement for Y_d turns out to be too strict. The reason is simply that, in order to accommodate both CKM mixing among quarks and MNS mixing among leptons, we need to set

$$Y_d = \Theta_L V_{CKM}^* Y_d^D \Theta_R U_{MNS} \Theta_\nu, \quad (6)$$

in the basis where Y_u and the right-handed neutrino mass matrix are diagonal (see below). $Y_d^D = \text{diag}(Y_d, Y_s, Y_b)$ is the positive, diagonalized Y_d matrix. Θ_L, Θ_R and Θ_ν are diagonal phase matrices. Because of this, we will instead

take[§]

$$W = \frac{1}{2}(Y_u)_{ij}16_i16_j10_u + \frac{1}{2}(Y_d)_{ij}16_i16_j\frac{\langle 45 \rangle}{M_{Pl}}10_d. \quad (7)$$

Because of the combination of the Higgs multiplet 45, whose VEV $\langle 45 \rangle \neq 0$ breaks $SO(10)$, and the Higgs in 10, the effective Yukawa coupling being either in 10 (symmetric between two 16's) or 120 (anti-symmetric between two 16's) representations, the matrix Y_d can now have a mixed symmetry. We imagine that $SO(10)$ is broken to $SU(5)$ around 10^{17} GeV, and this operator is large enough for the down-type Yukawa matrix. Note that we define V and U matrices to be in the CKM form with only one CP violating phase each. Phases in Θ_L and Θ_R are relevant only when the superheavy color triplet components of the Higgs multiplet is involved. Θ_ν is relevant for the CP violation in the neutrino sector when the Majorana character of the neutrino mass is involved.

Now we break $SO(10)$ to $SU(5)$. Given a strong hierarchy among up-type Yukawa couplings and the large top Yukawa coupling, it is natural to stick to the basis where Y_u is diagonal, $(Y_u)_{ij} = (Y_u^D)_i\delta_{ij}$. Further decomposing multiplets under $SU(5)$ as $16_i = 10_i + \bar{5}_i + 1_i$, and keeping only the Higgs multiplets $5_u \in 10_u$ and $\bar{5}_d \in 10_d$, we find ¶

$$W = \frac{1}{2}(Y_u^D)_i10_i10_i5_u + (Y_u^D)_i\bar{5}_i1_i5_u + (Y_d)_{ij}10_i\bar{5}_j\bar{5}_d + \frac{1}{2}M_{ij}1_i1_j. \quad (8)$$

It is clear that, in the absence of the right-handed neutrino mass matrix, we can eliminate U_{MNS} entirely by changing the basis of $\bar{5}_i$ in the $SU(5)$ superpotential.^{||} This is an immediate way to see that the only effect of U_{MNS} is related to the neutrino mass.

Further breaking $SU(5)$ down to the standard model, the prediction of this framework is that the Yukawa couplings in the MSSM+N (the MSSM together with right-handed neutrinos) are

$$W = (Y_u^D)_iQ_iU_iH_u + (Y_u^D)_iL_iN_iH_u + (V^*Y_d^D\Theta_R U\Theta_\nu)_{ij}Q_iD_jH_d + (V^*Y_d^D\Theta_R U\Theta_\nu)_{ij}E_iL_jH_d + \frac{1}{2}M_{ij}N_iN_j, \quad (9)$$

[§]The absence of renormalizable Yukawa coupling to 10_d could well be a consequence of discrete symmetries.

¶Right-handed neutrino masses arise from the coupling to $SO(10)$ -breaking Higgses either $\langle 126 \rangle$ or $\langle 16 \rangle^2/M_{Pl}$, as usual.

^{||}This, of course, is not necessarily true with the soft terms, which is the whole point of this paper.

where we had absorbed the phases in Θ_L into Q_i and E_i . Y_d is diagonalized by a bi-unitary rotation where the matrix acting on the left side represents the relative rotation of the left-handed down quarks with respect to the left-handed up quarks in the basis where the up quark mass matrix is diagonal. Hence, such matrix is just the usual CKM mixing matrix. The matrix acting on the right side represents the rotation to be performed on the left-handed leptons to go the mass basis of the charged leptons. It is easy to see that the phase matrix Θ_ν can be absorbed into the Majorana mass matrix M after redefining D_j and L_j , and, the phase matrix Θ_R can be absorbed into (UD) multiplet or (EV^*) multiplet. The phase matrices $\Theta_{L,R}$ are irrelevant as long as the colored triplet Higgs boson can be ignored as emphasized before. Hence such matrix U is to be identified with the neutrino mixing matrix U_{MNS} if we are in a basis where the physical light neutrinos are mass eigenstates. For this to happen, given that the neutrino Yukawa coupling matrix Y_u is diagonal, we have to assume that simultaneously also the right-handed neutrino mass matrix M is diagonal. Hence throughout our discussion we are taking Y_u and M *simultaneously diagonal*. Such a situation could result from simple $U(1)$ family symmetries. As we will comment below, in a scheme a la $SO(10)$ with hierarchical Y_u^D and right-handed neutrino masses, the choice of having such simultaneous diagonalization looks rather plausible.**

The similarity of the charged lepton and down-quark Yukawa matrices is well-known phenomenologically. Quantitatively, the relation $m_b = m_\tau$ could be indeed true at the unification scale, while $m_s = m_\mu$, $m_e = m_d$ are a factor of about three off. Here we take the point of view that the factors of three can be remedied by small $SU(5)$ -breaking effects of the framework and do not worry about it. Clearly, lower-generation Yukawa couplings are subject to more corrections simply because their sizes are small. The B -physics signatures we will discuss do not depend on such details as we end up ignoring all Yukawa couplings except that of the top quark. It is important to notice that the order of left- and right-handed fields is the opposite between $Q_i D_j$ and $E_i L_j$ couplings.

The important outcome of this framework is the (approximate) equality of the neutrino and up-quark Yukawa matrices. The light neutrino masses, after integrating out the right-handed neutrinos in Eq. (9), are given by the

**For a discussion of neutrino masses and mixings based on the simplest $SO(10)$ mass relations and the see-saw mechanism, see the work of Ref.[12]

superpotential

$$W = \frac{1}{2}(Y_u^D)_i(M^{-1})_{ij}(Y_u^D)_j(L_i H_u)(L_j H_u). \quad (10)$$

Since we assume that M is also diagonal in the same basis, this leads to the light Majorana neutrino mass matrix $(m_\nu)_{nm} = (Y_{ui}^{D2}/2M_i)U_{ni}^*e^{-i\delta_i}U_{mi}^*$, where $e^{i\delta_i}$ is the phase of M_i . The immediate conclusion is that the right-handed neutrino Yukawa matrix must be roughly doubly hierarchical compared to the up-quark Yukawa matrix. Phenomenologically, the Large Angle MSW solution is the most promising solution to the solar neutrino problem. Then the two mass splittings [13]

$$\Delta m_\oplus^2 \simeq 3 \times 10^{-3} \text{ eV}^2, \quad (11)$$

$$\Delta m_\odot^2 \simeq 0.3\text{--}2 \times 10^{-4} \text{ eV}^2, \quad (12)$$

are not very different, especially after taking their square root. On the other hand, the up-quark Yukawa matrix has a strong hierarchy $Y_u \ll Y_c \ll Y_t$. To obtain similar mass eigenvalues between the largest and the 2nd largest eigenvalues as suggested by data, we need $Y_c^2/M_2 \sim 0.2Y_t^2/M_3$. Moreover, the basis where the Y_u matrix is diagonal must be strongly correlated to the basis where the right-handed neutrino masses are diagonal to achieve this. The simplest possibility is to assume their simultaneous diagonalization, as we said above. Note also that all three physical CP violating phases associated with the light 3×3 Majorana mass matrix are present in this model as free parameters.

At GUT scale, the top quark mass and the largest neutrino Dirac mass are equal. As a result the heaviest neutrino mass is m_t^2/M_3 . From the recent fit to the Super-Kamiokande neutrino data and assuming non-degenerate neutrino masses, one has $m_{\nu_3} \sim 0.05 \text{ eV}$. For $m_t \sim 178 \text{ GeV}$, this corresponds to M_3 of roughly 10^{15} GeV slightly below the GUT scale as expected. It is very interesting to see that the $SO(10)$ model ties up neutrino mass, top mass and GUT scale nicely.

3.2 Effects on Soft Masses

The size of the radiative corrections on the SUSY soft masses induced by the neutrino Yukawa couplings and their possible consequence on low-energy flavor physics had been studied within the $SU(5)$ unification in Refs. [6,

14, 15, 16]. Following these papers, we shall assume that above some GUT unification scale, the SUSY breaking parameters are universal and can be parameterized by the universal scalar mass m_0 , the universal A -parameter a_0 which is the ratio of the SUSY breaking trilinear scalar interaction to the corresponding Yukawa couplings, the B -parameter entering the scalar bilinear term mixing the two Higgs doublets and the universal gaugino mass m_G . The scale, M_* , where these universal SUSY breaking values should be applied depends on the details of the SUSY breaking mechanism. Here we shall simply assume it to be near the Planck scale .

In the context of SUSY $SU(5)$ [6, 14], it was shown that, if the right-handed neutrino singlet is introduced to account for the data on neutrino oscillation, large neutrino Yukawa couplings involved in the neutrino Dirac masses can induce large off-diagonal mixings in the right-handed down squark mass matrix through renormalization group evolution between M_* and M_{GUT} . In addition, the contributions to the scalar masses induced in the running by the neutrino Yukawa couplings will generally be complex with new CP violating phases unrelated to the Kobayashi-Maskawa(KM) phase in Standard Model. The abovementioned mixings can be parameterized as $\delta_{ij}^R = (m_{\tilde{d}_R}^2)_{ij}/m_{\tilde{q}^2}$ where $m_{\tilde{q}^2}$ is the average right-handed down squark mass. In particular, in Ref. [6], it was shown that the induced δ_{12}^R is large enough to account for many of the observed CP violating phenomena in the kaon system providing an alternative to the CKM interpretation of these data. In Ref. [14], it was shown that δ_{13}^R can give rise to a CP asymmetry in $B_d \rightarrow \phi K_s$ much larger than the KM prediction.

Here we wish to point out first that the off-diagonal mixing parameter δ_{23} is further enhanced in the context of a SUSY $SO(10)$ model. In the next section we will elaborate on the phenomenological consequence of large δ_{23}^R . In our case the constraint coming from the upper bound on $BR(\mu \rightarrow e\gamma)$ turns out to be less severe than in the $SU(5)$ context.

Due to the larger matter content of the $SO(10)$ GUT model, the renormalization group evolution from M_* down to $SO(10)$ breaking scale, M_{10} , is faster than that of the $SU(5)$ model. The induced off-diagonal elements in the SUSY breaking mass matrix of the right-handed down squarks $\tilde{d}_R$ are given by (in the basis in which Y_D is diagonal)

$$[m_{\tilde{d}_R}^2]_{nm} \simeq -\frac{1}{8\pi^2} [Y^{u\dagger} Y^u]_{nm} (3m_0^2 + a_0^2) \left(5 \log \frac{M_*}{M_{10}} + \log \frac{M_{10}}{M_5} \right) \quad (13)$$

where M_5 is the $SU(5)$ breaking scale and

$$[Y^{u\dagger}Y^u]_{nm} = [(\Theta_R U Y_u^{D2} U^\dagger \Theta_R^*)]_{nm} = e^{-i(\phi_m^{(L)} - \phi_n^{(L)})} y_t^2 [U]_{m3}^* [U]_{n3}, \quad (14)$$

where $e^{i\phi_n^{(L)}}$ is the phase from $(\Theta_R)_{nn}$. Note that these phases are not relevant to any other low energy physics.

To account for the large atmospheric neutrino mixing, the second and third entries of the third row of U_{MNS} should be of order $\sqrt{2}$, while the first entry, U_{3e} , is severely limited by the CHOOZ experiment: $|U_{e3}| \leq 0.11$. Hence we obtain:

$$[Y^{u\dagger}Y^u]_{23} = 0.5e^{-i(\phi_2^{(L)} - \phi_3^{(L)})} (m_{tG}/178 \text{ GeV})^2, \quad (15)$$

where m_{tG} is the top quark mass at M_G .

The factor 5 in the RG coefficient above the $SO(10)$ breaking scale is due to the contribution of the loop diagram with $(10, \bar{5}_u)$ multiplets of $SU(5)$ in the loop which is not present in $SU(5)$. They contribute four times more than the usual $(1, 5_u)$ contribution in $SU(5)$, and δ_{23} can easily be $O(1)$.

Note that m_{tG} can be quite different from the pole mass of about $m_t \sim 178 \text{ GeV}$. The evolution of m_{tG} between M_* and M_G has been discussed in the literature [17].

In $SU(5)$ models, people had assumed that the right-handed neutrino mass matrix is given by an identity matrix to simplify the analysis. Given only a small hierarchy between $\Delta m_\oplus^2$ and $\Delta m_\odot^2$ for Large Mixing Angle MSW Solution, the second-generation neutrino Yukawa coupling is sizable, with a large mixing with the first-generation states. This led to quite stringent constraints from the processes such as $\mu \rightarrow e\gamma$. In our $SO(10)$ framework, however, the neutrino Yukawa matrix is as hierarchical as the up-quark sector, and only the third-generation Yukawa coupling is significant. In the third-generation multiplet, the electron state appears with a suppressed coefficient U_{e3} . Therefore, unlike the frameworks studied in the literature, contributions to processes that involve the first generation (like $\mu \rightarrow e + \gamma$) are suppressed by this unknown element in the MNS matrix. In view of this fact, we prefer to focus on flavor violating processes involving the mixing between second- and third-generation in this paper.




Figure 1: Possible important contributions to B -physics from large $\tilde{b}_R$ - $\tilde{s}_R$ mixing, such as B_s - $\bar{B}_s$ mixing, and SUSY penguin contribution to $B_d \rightarrow \phi K_s$ transition.

4 Consequences in B Physics

In this Section we present some implications of a large and complex δ_{23}^R in B physics. The discussion is semi-quantitative. Fully quantitative evaluation of effects in B -physics and their correlations will be discussed elsewhere.^{††}

The diagrammatic contributions of δ_{23}^R to various $\Delta B = -\Delta S = 2$ and $\Delta B = -\Delta S = 1$ processes were worked out in detail in Ref. [18]^{‡‡}. In particular a complex δ_{23}^R can play a major role in CP violating B decays [20, 14, 21, 16, 22].

The first effect of a conspicuous δ_{23}^R would be a large contribution to the $\Delta B = -\Delta S = 2$, B_s - $\bar{B}_s$ mixing through the operator $Q_1 = \bar{s}_R^\alpha \gamma_\mu b_R^\alpha \bar{s}_R^\beta \gamma^\mu s_R^\beta$ with complex coefficient

$$\mathcal{H}_{eff} = -\frac{\alpha_s^2}{216m_{\tilde{q}}^2} (\delta_{23}^d)_{RR}^2 \left(24 Q_1 x f_6(x) + 66 Q_1 \tilde{f}_6(x) \right) \quad (16)$$

where the functions f_6 and $\tilde{f}_6$ are defined as in Ref. [18].

From Eqs. (13,15) we see that δ_{23}^R can easily be as large as 0.5, yielding a SUSY contribution to ΔM_s comparable to that of the SM. Hence, in our scheme the operator gives rise to a large B_s - $\bar{B}_s$ mixing with a complex phase

^{††}We stick to mass insertion formalism for illustrative purposes, but fully quantitative discussions would require calculations in the mass eigenbasis.

^{‡‡}For an updated analysis of the gluino-mediated SUSY contributions to the B_d - $\bar{B}_d$ mixing and to the CP asymmetry in the decay $B \rightarrow J/\psi K_s$ including the NLO QCD corrections and B coefficients as computed in the lattice instead of using the vacuum insertion approximation, see Ref. [19]

which is almost unconstrained so far. In contrast, the SM contribution to B_s - $\bar{B}_s$ mixing has very small phase (in the usual Wolfenstein convention). This gives rise to many phenomenological consequences as will be sampled below.

Secondly, δ_{23}^R gives rise to new contribution to direct B decays. Two categories of contributions are more important. The first one is a $\Delta B = 1$ box diagram contribution. The second is of the type of electromagnetic or gluonic Penguin contributions.* There are also contribution of electroweak Penguin type which we shall ignore because they are generally smaller.

In B decay processes in which the dominant contribution in SM is at the tree level, the additional contribution due to δ_{23}^R can at most be a small percentage. This is true even if the SM contribution comes with strong mixing angle suppression [22] such as $B^\pm \rightarrow DK^\pm$ or $\bar{D}K^\pm$. However, if the initial state mixing (such as B_S mixing) plays a strong role in the phenomena, then δ_{23}^R can significantly alter the phenomenology.

In B decay processes in which the dominant contributions involve one loop contributions, such as the Penguin diagrams, one should expect large additional contribution due to the δ_{23}^R in both amplitude and CP asymmetry.

As an application of the above analysis, we can roughly classify the phenomenology into three categories.

- (a). Measurements of the β angle of the unitarity triangle. The leading mode, $B \rightarrow J/\psi K_s$, has large phase from the initial state B_d mixing (in Wolfenstein convention), and large real tree-level decay amplitude. It therefore does not receive significant contribution from δ_{23}^R . However, other modes, such as $B_d \rightarrow \phi K_s$ ($\bar{b} \rightarrow \bar{s}s\bar{s}$), which, in SM, measures the same β , can now receive large additional contribution from δ_{23}^R through the box and Penguin diagrams [14, 21]. The fractional phase difference $r_\beta = (\beta(JK_s) - \beta(\phi K_s))/\beta(JK_s)$ is a measure of δ_{23}^R that can be as large as 50%.
- (b). Measurements of the γ angle of the unitarity triangle. One class of popular measurements on γ involves B_s decays [24]. In the SM, the B_s mixing is real to a very good approximation. Therefore any measurements of γ using B_s decays will be strongly affected by δ_{23}^R . For

*Fully quantitative analysis would require a careful study of possible cancellation between penguin and box diagrams [23].

example, in $\bar{B}_s \rightarrow (D_s)^- K^+$ the CP asymmetry is due to the interference of the B_s decays, which has real amplitude in SM, and the $\bar{B}_s$ decay with complex amplitude after the B_s - $\bar{B}_s$ mixing. The two decays are roughly of equal magnitude and the phase of $\bar{B}_s$ is exactly γ in SM. With δ_{23}^R , even the $\bar{B}_s \rightarrow B_s \rightarrow (D_s)^- K^+$ develops a large phase due to additional complex contribution to the mixing. The phase is proportional to $\text{Arg}(M_{12}(\delta_{23}^R)/(M_{12}(\delta_{23}^R) + M_{12}(\text{SM})))$ where $M_{12}(\text{SM})$ and $M_{12}(\delta_{23}^R)$ are the B_s mixing amplitude of the SM and that due to δ_{23}^R respectively [22].

Another class of measurement on γ are using charge B decays. For example in $B^+ \rightarrow D^0 K^+$ or $\bar{D}^0 K^+$, γ is measured through the interference between the two quark level processes $b \rightarrow c\bar{u}s$ and $b \rightarrow u\bar{c}s$. The decay chains of c or $\bar{c}$ in the final state produce a D^0 or $\bar{D}^0$ mesons respectively. The two contributions interfere if both D^0 or $\bar{D}^0$ decay to the same final state f_D and have a relative phase γ . Here, f_D is one of the states that both D and $\bar{D}$ can decay into, such as $K^-\pi^+$ or CP eigenstates K^+K^- , $\pi^+\pi^-$, $K_s\pi^0$ or $K_s\phi$ [22]. In this measurement, the role played by δ_{23}^R is negligible, so it measures the same value as in SM. Therefore, just like β measurements, by comparing γ measurements in B_s and in $B^\pm$ decays, one can get a measure of δ_{23}^R . In principle, by comparing r_β with r_γ , which is similarly defined, one can get a strong evidence of the existence of large δ_{23}^R .

- (c). Decays which are expected to be essentially CP conserving in the SM. Some of decays may have large CP asymmetry due to the existence of δ_{23}^R . Examples: $B_s \rightarrow J\phi$ or $B_s \rightarrow (D_s)^+(D_s)^-$ or $B \rightarrow X_s\gamma$.

5 Conclusion

In this Letter, we pointed out that a large mixing between ν_τ and ν_μ as observed in atmospheric neutrino oscillation may lead to a large mixing between $\tilde{b}_R$ and $\tilde{s}_R$ because they belong to the same $SU(5)$ multiplets. This occurs naturally in $SO(10)$ grand unified models which we have described in detail. These models do not give rise to dangerously large $\mu \rightarrow e\gamma$ and similar processes which involve the first generation, given the current limit on U_{e3} from reactor-neutrino experiments. A large mixing between $\tilde{b}_R$ and $\tilde{s}_R$ leads to interesting effects in B -physics, such as large and CP-violating

B_s mixing, different “ $\sin 2\beta$ ” between $B_d \rightarrow \phi K_s$ and $J/\psi K_s$, different “ γ ” from various measurements, and CP asymmetry in $B_s \rightarrow J\phi$, $(D_s)^+(D_s)^-$.

Acknowledgements.

HM was supported in part by the Director, Office of Science, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-0098840. DC is supported by a grant from NSC of ROC and LBL. DC wishes to thank the Theory Group at LBL and the Theory Group at SLAC for hospitality while this work is in progress and Lincoln Wolfenstein for discussion. AM thanks the UCB and LBL Theory Groups for kind hospitality and support at the early stages of this work.

Note Added

After this manuscript is submitted to the arXiv, we learned from Jogesh Pati that a similar idea by K.S. Babu and J. Pati was mentioned in his talk at the CasCais, Portugal School (July 2000), the abstract of which appeared in the Proceedings edited by Branco, Shafi and Silva-Marcos, Pages 215.

References

- [1] A. Masiero and H. Murayama, Phys. Rev. Lett. **83**, 907 (1999) [arXiv:hep-ph/9903363].
- [2] T. Yanagida, in *Proc. of the Workshop on the Unified Theory and Baryon Number in the Universe*, ed. O. Sawada and A. Sugamoto (KEK report 79-18, 1979), p. 95;
M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, ed. P. van Nieuwenhuizen and D.Z. Freedman (North Holland, Amsterdam, 1979), p. 315.
- [3] E. Witten, Phys. Lett. B **91**, 81 (1980);
R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980);
T. Yanagida, Prog. Theor. Phys. **64**, 1103 (1980).
- [4] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57** 961 (1986)

- [5] W. Buchmuller, D. Delepine and F. Vissani, Phys. Lett. **B459** 171 (1999); W. Buchmuller, D. Delepine and L. T. Handoko, Nucl. Phys. **B576** 445 (2000); J. Ellis, M. E. Gomez, G. K. Leontaris, S. Lola and D. V. Nanopoulos, Eur. Phys. J. **C14** 319 (2000); J. Hisano and K. Tobe, Phys. Lett. **B510** 197 (2001); J. A. Casas and A. Ibarra, arXiv:hep-ph/0103065; D. F. Carvalho, J. Ellis, M. E. Gomez and S. Lola, arXiv:hep-ph/0103256; T. Blazek and S. F. King, arXiv:hep-ph/0105005; J. Sato, K. Tobe and T. Yanagida, Phys. Lett. **B498** 189 (2001); J. Sato and K. Tobe, Phys. Rev. **D63** 116010 (2001); S. Lavignac, I. Masina and C. A. Savoy, arXiv:hep-ph/0106245.
- [6] T. Moroi, JHEP. **0003** 019 (2000), [arXiv:hep-ph/0002208]; N. Akama, Y. Kiyo, S. Komine and T. Moroi, Phys. Rev. **D64** 095012 (2001), [arXiv:hep-ph/0104263].
- [7] L. Alvarez-Gaume, M. Claudson and M. B. Wise, Nucl. Phys. B **207**, 96 (1982);
M. Dine and W. Fischler, Phys. Lett. B **110**, 227 (1982);
M. Dine and A. E. Nelson, Phys. Rev. D **48**, 1277 (1993) [arXiv:hep-ph/9303230];
M. Dine, A. E. Nelson and Y. Shirman, Phys. Rev. D **51**, 1362 (1995) [arXiv:hep-ph/9408384].
- [8] L. Randall and R. Sundrum, Nucl. Phys. B **557**, 79 (1999) [arXiv:hep-th/9810155];
G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP **9812**, 027 (1998) [arXiv:hep-ph/9810442].
- [9] H. Georgi, in *Particles and Fields*, ed. C. E. Carlson (AIP, NY, 1975), p.575; H. Fritzsch and P. Minkowski Ann.of Phys. **93** 193 (1975).
- [10] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962).
- [11] S. Dimopoulos and L. J. Hall, Phys. Lett. **B344**, 185 (1995) [arXiv:hep-ph/9411273].
- [12] W. Buchmueller and D. Wyler, Phys. Lett. **B521**, 291 (2001) [arXiv:hep-ph/0108216]
- [13] Q. R. Ahmad *et al.* [SNO Collaboration], arXiv:nucl-ex/0204009.

- [14] T. Moroi, Phys. Lett. **B493**, 366 (2000) [arXiv: hep-ph/0007328].
- [15] P. Ciafaloni, A. Romanino and A. Strumia Nucl. Phys. **B458**, 3 (1996) , [arXiv:hep-ph/9507379]; J. Hisano, T. Moroi, K. Tobe, M. Yamaguchi, T. Yanagida, Phys. Lett. **B357**, 579 (1995), [arXiv:hep-ph/9501407] ; J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. **D53**, 2442 (1996), [arXiv:hep-ph/9510309]; J. Hisano, T. Moroi, K. Tobe, M. Yamaguchi, Phys. Lett. **B391**, 341 (1997), Erratum-ibid. **B397** 357 (1997), [arXiv:hep-ph/9605296].
- [16] R. Barbieri, L. J. Hall and A. Strumia, Nucl. Phys. **B449**, 437 (1995), [arXiv:hep-ph/9504373] ; M. B. Causse, J. Orloff, [arXiv:hep-ph/0012113]; A. Masiero, O. Vives, Nucl. Phys. Proc. Suppl. **99B**, 228 (2001), [arXiv:hep-ph/0107276]; Phys. Rev. Lett. **86**, 26 (2001), [arXiv:hep-ph/0007320]; D. A. Demir, A. Masiero and O. Vives, Phys. Rev. Lett. **82**, 2447 (1999), [arXiv:hep-ph/9812337]; G. Barenboim, K. Huitu and M. Raidal Phys. Rev. **D63** 055006 (2001), [arXiv:hep-ph/0005159]; S. Baek, T. Goto, Y. Okada and K. Okumura Phys. Rev. **D63** 051701 (2001), [arXiv:hep-ph/0002141]; Phys. Rev. **D64** 095001 (2001), [arXiv:hep-ph/0009196]; T. Goto, Y. Okada, Y. Shimizu, T. Shindou and M. Tanaka, arXiv:hep-ph/0204081.
- [17] R. Barbieri, L. J. Hall and A. Strumia, Nucl. Phys. **B445** 219 (1995), [arXiv:hep-ph/9501334].
- [18] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, Nucl. Phys. **B477** 321 (1996), [arXiv:hep-ph/9604387].
- [19] D. Becirevic *et al.*, arXiv:hep-ph/0112303.
- [20] M. Ciuchini, E. Franco, G. Martinelli, A. Masiero and L. Silvestrini, Phys. Rev. Lett. **79** 978 (1997) [arXiv:hep-ph/9704274].
- [21] E. Lunghi and D. Wyler, Phys. Lett. **B521** 320 (2001), [arXiv:hep-ph/0109149].
- [22] D. Chang, W.-F. Chang, W.-Y. Keung, N. Sinha and R. Sinha, Phys. Rev. **D65** 055010 (2002), [arXiv:hep-ph/0109151].
- [23] Y. Grossman, M. Neubert and A. L. Kagan, JHEP **9910**, 029 (1999) [arXiv:hep-ph/9909297].

[24] I. I. Bigi and A. I. Soni, in *CP Violation*, Cambridge Univ. Press. (2000).